

Apolipoprotein L1 High-Risk Genotypes and Albuminuria in Sub-Saharan African Populations

Jean-Tristan Brandenburg ¹, Melanie A. Govender ^{1,2}, Cheryl A. Winkler ³, Palwende Romuald Boua ^{1,4}, Godfred Agongo ^{5,6}, June Fabian ^{7,8} and Michèle Ramsay ^{1,2}

Abstract

Background and objectives Recessive inheritance of African-specific *APOL1* kidney risk variants is associated with higher risk of nondiabetic kidney disease, progression to kidney failure, and early-onset albuminuria that precedes eGFR decline. The effect of *APOL1* risk variants on kidney disease in continental Africans is understudied. Objectives of this study were to determine *APOL1* risk allele prevalence and associations between *APOL1* genotypes and kidney disease in West, East, and South Africa.

Design, setting, participants, & measurements This cross-sectional population-based study in four African countries included 10,769 participants largely aged 40–60 years with sociodemographic and health information, anthropometry data, and blood and urine tests for biomarkers of kidney disease. *APOL1* risk alleles were imputed from the H3Africa genotyping array, *APOL1* risk allele and genotype frequencies were determined, and genetic associations were assessed for kidney disease. Kidney disease was defined as the presence of eGFR <60 ml/min per 1.73 m², albuminuria, or a composite end point including eGFR <60 ml/min per 1.73 m² and/or albuminuria.

Results High G1 allele frequencies occurred in South and West Africa (approximately 7%–13%). G2 allele frequencies were highest in South Africa (15%–24%), followed by West Africa (9%–12%). Associations between *APOL1* risk variants and albuminuria were significant for recessive (odds ratio, 1.63; 95% confidence interval, 1.25 to 2.12) and additive (odds ratio, 1.39; 95% confidence interval, 1.09 to 1.76) models. Associations were stronger for *APOL1* G1/G1 genotypes versus G0/G0 (odds ratio, 3.87; 95% confidence interval, 2.16 to 6.93) compared with either G2/G2 (odds ratio, 1.65; 95% confidence interval, 1.09 to 2.51) or G1/G2 (odds ratio, 1.24; 95% confidence interval, 0.83 to 1.87). No association between *APOL1* risk variants and eGFR <60 ml/min per 1.73 m² was observed.

Conclusions *APOL1* G1 and G2 alleles and high-risk genotype frequencies differed between and within West and South Africa and were almost absent from East Africa. *APOL1* risk variants were associated with albuminuria but not eGFR <60 ml/min per 1.73 m². There may be differential effects of homozygous G1 and G2 genotypes on albuminuria that require further investigation.

CJASN 17: ●●–●●, 2022. doi: <https://doi.org/10.2215/CJN.14321121>

Introduction

Prevalent CKD is estimated at 8%–16% worldwide, with high morbidity and mortality in resource-limited countries (1). The true prevalence of CKD and phenotypic associations with genetic risk factors are understudied in resident African populations (2). By contrast, CKD and associated genetic risk are well characterized in admixed African American populations, informed by a series of landmark studies. These studies showed that focal segmental glomerulosclerosis (FSGS), hypertension-attributed kidney disease, and human immunodeficiency virus-associated nephropathy (HIVAN) were strongly associated with three African-specific apolipoprotein L1 (*APOL1*) coding variants on chromosome 22 (3–8).

The chromosome 22 region was discovered by admixture mapping and found to harbor African-specific

kidney risk variants that include two missense single nucleotide polymorphisms (SNPs) defining the G1 haplotype (rs73885319 and rs60910145) and a six-base-pair deletion removing two amino acids (rs71785313) defining the G2 haplotype (9). Most likely, G1 and G2 variants originated in West Africa, and their frequency increased due to recent positive selection by *Trypanosoma brucei rhodesiense* in East Africa and *Trypanosoma brucei gambiense* in West Africa (10).

Subsequent studies showed that *APOL1* risk alleles were not associated with diabetic or IgA nephropathy but were associated with collapsing glomerulopathies due to systemic lupus erythematosus, exogenous interferon (IFN) administration, untreated HIV infection, and, more recently, severe acute respiratory syndrome coronavirus 2 (6,11–15). To date, the

Due to the number of contributing authors, the affiliations are listed at the end of this article.

Correspondence:

Dr. Jean-Tristan Brandenburg or Prof. Michèle Ramsay, 9 Jubilee Road, Parktown 2193, South Africa. Email: jean-tristan.brandenburg@wits.ac.za or michele.ramsay@wits.ac.za

spectrum of *APOL1*-associated disease includes sickle cell nephropathy, preeclampsia, FSGS, and hypertension-attributed kidney disease (16). In the African American population, *APOL1* allele frequencies are approximately 23% for G1 and 13% for G2 (6). Almost 50% of the African American population carries one risk allele, and approximately 13% of the African American population carries two risk alleles (G1/G1, G1/G2, and G2/G2), comprising the high-risk genotype for kidney disease (17). Although case-control studies contribute to understanding *APOL1*-associated kidney disease (often reporting high odds ratios [ORs]), these are limited by selection and survival biases and require corroboration in population-based studies (18).

Large community-based studies have confirmed higher risk for incident kidney disease (albuminuria; eGFR < 60 ml/min per 1.73 m²), progression to kidney failure, and early-onset albuminuria that precedes decline in eGFR in African American populations with *APOL1* high-risk genotypes compared with other US population groups (7,19–23). Most studies support a recessive model of inheritance with little or no differential effects between high-risk genotypes (5,6,11).

However, differential effects between G1 and G2 homozygotes have been reported. When compared with G2 homozygotes, G1 homozygotes in the Dallas Heart study had a higher incidence of kidney disease (eGFR < 60 ml/min per 1.73 m²) overall ($P=0.05$) and for nondiabetic kidney disease ($P=0.007$), and G1 homozygotes in the Jackson Heart study had higher risks for CKD ($P=0.04$) and albuminuria ($P=0.02$) (7,24). Furthermore, G1/G0 heterozygosity has been associated with higher risk for hypertension-associated kidney failure, younger age at presentation of nondiabetic kidney failure, and higher risk for FSGS or HIVAN, whereas G2/G0 heterozygosity is present in all documented cases of IFN-associated nephropathy (5,6,15,25–27).

Although frequency distributions of *APOL1* alleles have been reported, no population-based studies and only a handful of case-control studies have investigated the spectrum of *APOL1*-associated kidney disease in sub-Saharan Africa. Regarding *APOL1* allele frequencies, some West African ethnic groups from Ghana and Nigeria have G1 frequencies >40% with lower G2 frequencies (6%–24%). Previously published studies reported lower combined G1 and G2 frequencies (approximately 18.4%) in South Africa, and absent or infrequent occurrence in East Africa (15,17,28).

Of the available patient-control studies, *APOL1* high-risk genotypes have been associated with hypertension-attributed kidney disease in the Democratic Republic of Congo, specifically G1/G1 and G1/G2 (OR, 7.7; 95% confidence interval [95% CI], 1.5 to 39.7) (29); nondiabetic kidney disease in Nigeria (OR, 4.8; 95% CI, 1.6 to 14.9) (30); and HIVAN in South Africa (OR, 89; 95% CI, 18 to 912) (15). The *APOL1* association with HIVAN differs remarkably between South Africans (OR, 89; 95% CI, 18 to 912) and African American participants (OR, 29; 95% CI, 13.1 to 68.5), although approximately the same number (78% and 72%, respectively) of cases carried high-risk genotypes (6,15). In the South African study (but not the US study), the G1/G0 genotype was strongly associated with HIVAN risk. In a Nigerian study of HIV-infected participants on antiretroviral therapy, *APOL1* high-risk genotypes were

associated with a higher likelihood of microalbuminuria (OR, 1.97; 95% CI, 1.37 to 2.82) and macroalbuminuria (OR, 3.96; 95% CI, 1.95 to 8.02); this was the first study from the region to investigate the effect of *APOL1* high-risk genotypes on early kidney disease (31).

In our large population-based study from West, East, and South Africa, we performed an exploratory analysis to characterize *APOL1* genetic variation, compare allele and genotype frequencies within and between study sites, and investigate associations between *APOL1* genotypes and biomarkers of kidney disease: low eGFR, albuminuria, and a composite end point comprising low eGFR and/or albuminuria.

Materials and Methods

Study Participants

This study was part of the Africa Wits–International Network for the Demographic Evaluation of Populations and Their Health Partnership for Genomic Studies (referred to as AWI-Gen), a cross-sectional population-based study of over 12,000 participants largely aged 40–60 years at recruitment from six participating urban and rural centers in four sub-Saharan African countries. Pregnant women, first-degree relatives of existing participants, recent immigrants (with <10 years of residence in the region), and individuals with physical impairments preventing measurement of BP and other anthropometric indices were excluded. West Africa included two countries: Ghana (Navrongo) and Burkina Faso (Nanoro), East Africa included Kenya (Nairobi), and South Africa had three study sites (Soweto, Agincourt, and Dikgale). The overall aim of the AWI-Gen collaboration is to investigate genomic and environmental risk factors that affect cardiometabolic disease (32,33).

For this study, 10,769 adults with data for *APOL1* genotypes were included. The Human Research Ethics Committee of the University of the Witwatersrand, South Africa, approved the study, with additional institutional review board approval from each participating site. All participants provided written informed consent, and the study was conducted in accordance with the Declaration of Helsinki. Study methods are detailed elsewhere (33), but briefly, cross-sectional demographic, health, and family history information was obtained through a researcher-administered questionnaire. Self-reported ethnicity was collected to assess diversity of participants. Anthropometric measurements, clinical data, fasting venous blood samples, and spot midstream urine samples were collected.

This study received the approval of the Human Research Ethics Committee (Medical), University of the Witwatersrand/South Africa (M121029 and M170880); the Centre Muraz Institutional Ethics Committee/Burkina Faso (015-2014/CE-CM); the National Ethics Committee for Health Research/Burkina Faso (2014-08-096); the Ghana Health Service Ethics Review Committee (identification no. GHS-ERC:05/05/2015); the Navrongo Institutional Review Board (identification no. NHRCIRB178); and the AMREF Health Ethics and Scientific Review Committee in Kenya (approval no. P114/2014). All of the participants provided informed consent before entering the study.

Biomarkers of Kidney Disease

Biomarkers of kidney disease were defined as low eGFR (eGFR <60 ml/min per 1.73 m²), albuminuria (random spot urine: urine albumin-creatinine ratio [UACR] >30 mg/g), and a composite end point including low eGFR and/or albuminuria. The Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI; creatinine) 2009 equation without adjustment for African American ethnicity (34) was used to estimate the GFR.

At each partner site, blood and urine specimens were processed and stored at -80°C. After completion of the study, each partner site transported samples on dry ice to a central laboratory in Johannesburg, South Africa. At the central laboratory, all analyses were batched and performed according to good laboratory practice with external monitoring for quality control. Serum creatinine and urine creatinine (milligrams per deciliter) were measured using the Jaffe kinetic method calibrated to an isotope dilution mass spectrometry-traceable standard. Urinary albumin concentration was measured with immunoturbidimetry. The UACR (milligrams per gram) was calculated from these measurements. DNA extracted from buffy coat samples was used for genotyping.

Comorbidities and Socioeconomic Status

Diabetes mellitus was defined as fasting blood glucose \geq 126 mg/dl, random blood glucose >198 mg/dl, self-reported history of diabetes mellitus, or currently on treatment regardless of the glucose concentration. Hypertension was defined as mean systolic BP \geq 140 mm Hg, mean diastolic BP \geq 90 mm Hg, self-reported history of hypertension, or currently on treatment regardless of BP measurement. HIV status was self-reported or diagnosed with a rapid test. Socioeconomic status was assessed using a validated tool appropriate for assessing wealth in low- and middle-income settings. Household assets are scored, and on the basis of the score, data are represented as quintiles, with the poorest in the first quintile (33).

Genotyping and Imputation of *APOL1* Genotypes

APOL1 G1 and G2 are formally haplotypes (Supplemental Table 1), but for ease of understanding, we refer to them as alleles in this publication. The *APOL1* haplotypes carrying neither G1 nor G2 variants are termed G0. *APOL1* high-risk genotypes comprise two risk alleles (G1/G1, G1/G2, and G2/G2), and low-risk genotypes comprise zero or one risk allele (G0/G0, G0/G1, and G0/G2).

Genomic DNA was genotyped using the H3Africa genotyping array (<https://www.h3abionet.org/h3africa-chip>) designed as an African common variant-enriched GWAS array (Illumina) with approximately 2.3 million SNPs. Preimputation quality control was applied to the entire AWI-Gen genotype dataset. Individuals with a missing SNP calling rate >0.05 were removed. SNPs with genotype missingness >0.05, minor allele frequency <0.01, and Hardy-Weinberg equilibrium *P* value <0.001 were removed. Nonautosomal and mitochondrial SNPs and ambiguous SNPs that did not match the GRCh37 reference alleles or strands were removed.

Imputation was performed on the cleaned dataset (with 1,729,661 SNPs and 10,903 individuals) with the UK10K and 1000 Genomes Project panel, using the Sanger imputation service (35). We extracted 187 positions between

35649056 and 37663576 on chromosome 22 around the *APOL1* gene from AWI-Gen GWAS data and used three whole-genome sequencing datasets to assess imputation accuracy: AWI-Gen whole-genome sequencing (36), 1000 Genomes Project (37), and the African Genome Variation Project (38). Three genomics positions were used to define the *APOL1* risk alleles G1 and G2: rs73885319 (p.S342G), rs60910145 (p.I384M), and insertion/deletion rs71785313 (p.N388Y389/-) (Supplemental Table 1). The imputation of these three variants was compared with the original sequences from the whole-genome sequencing datasets.

Statistical Analyses: Genetic Association with Biomarkers of Kidney Disease

We used mixed effects models using the ImerTest package (39) to compute associations between *APOL1* alleles and biomarkers of kidney disease. Logistic mixed models were used for binary biomarkers (low eGFR, albuminuria, and the composite end point for kidney disease). For analyses stratified by site, we used logistic models for binary biomarkers. ORs with 95% CIs were calculated for the association between biomarkers of kidney disease and (1) zero or one versus two *APOL1* risk alleles (recessive model), (2) zero versus one or two *APOL1* risk alleles (dominant model), or (3) three genotype groups (zero versus one versus two risk alleles; additive model). The models were adjusted for age, sex, body mass index (BMI), diabetes mellitus, hypertension, and HIV status as fixed variables and study site as a random variable. ORs and statistical significance were assessed using mixed effects models for carriage of one or two *APOL1* risk alleles in the recessive, dominant, and additive models. *APOL1* analysis was stratified by diabetes mellitus, hypertension, and HIV status, and mixed effects models were used to assess associations with albuminuria in recessive models. Furthermore, GxG interaction terms were computed independently using interaction terms between *APOL1* and diabetes mellitus, hypertension, and HIV in models adjusted for covariates.

For tables of descriptive data, the proportion of missing data for each variable was indicated using the appropriate denominator (different from the stipulated sample size) or annotated as a footnote. Missing data were not imputed. For UACR, data were missing at random for the Agincourt and Soweto sites, South Africa. Of note, there were no UACR data for women participants in Soweto. To assess potential effects of these missing data on the association between high-risk *APOL1* genotypes and albuminuria, fixed effects meta-analyses using results from linear mixed models by site and by sex confirmed that no significant bias was introduced from the missing data. To assess whether there was a site effect for the association between high-risk *APOL1* genotypes and albuminuria, we tested for heterogeneity using the Cochrane test.

Results

Genotype Imputation Quality

The rs73885319 SNP defining the G1 allele was present in the H3Africa array, and there was full concordance between the genotyped allele calls and imputed calls. Imputation accuracy values using African samples for rs60910145 (in linkage disequilibrium with the functional

G1 rs73885319 SNP) and rs71785313 (G2) were 99.9% and 99.0%, respectively, between whole-genome sequence and imputed data. Similarly, other populations showed concordance between G0, G1, and G2 alleles and imputed data assessed using sequence data (Supplemental Table 2).

Population Characteristics and *APOL1* Allele and Genotype Frequencies in South, East, and West African AWI-Gen Studies Sites

Demographic and clinical characteristics and *APOL1* allele and genotype frequency distributions for each study

Table 1. Population characteristics and *APOL1* allele and genotype frequencies in the Africa Wits–International Network for the Demographic Evaluation of Populations and Their Health Partnership for Genomic Studies

Variable ^a	Africa Wits–International Network for the Demographic Evaluation of Populations and Their Health Partnership for Genomic Studies Site						
	South Africa			East Africa	West Africa		Overall
	Agincourt, Rural	Soweto, Urban	Dikgale, Rural	Nairobi, Urban	Nanoro, Rural	Navrongo, Rural	All Sites
Sample size, <i>N</i>	2253	1775	1143	1771	1983	1844	10,769
Clinical phenotype							
Age, yr	58 (11)	49 (6)	52 (8)	49(6)	50(6)	51 (6)	52 (8)
Men, <i>N</i> (%)	938 (42)	922 (52)	361 (32)	804 (45)	1009 (51)	842 (46)	4876 (45)
Women, <i>N</i> (%)	1315 (58)	853 (48)	782 (68)	967 (55)	974 (49)	1002 (64)	5893 (55)
Height, m	1.65 (0.85)	1.65 (0.88)	1.62 (0.84)	1.64 (0.87)	1.68 (0.87)	1.62 (0.86)	1.65 (0.89)
Weight, kg	74 (17)	78 (19)	72 (20)	68 (14)	59 (12)	57 (10)	68 (17)
BMI, kg/m ^{2b}	27.2 (6.7)	28.9 (7.7)	27.8 (8.2)	25.4 (5.7)	21.0 (3.5)	21.6 (3.6)	25.1 (6.7)
Diabetes mellitus, <i>N</i> (%) ^c	194/2214 (9)	167/1770 (9)	113/1136 (10)	124/1755 (7)	72/1973 (4)	25/1794 (1)	695/10,642 (7)
Hypertension, <i>N</i> (%)	1265/2253 (56)	949/1775 (54)	515/1143 (45)	486/1771 (27)	322/1983 (16)	455/1844 (25)	3992/10,769 (37)
HIV infection, <i>N</i> (%) ^d	602/2251 (27)	253/1207 (21)	224/1124 (20)	216/1561 (14)	8/1983 (0.4)	15/1844 (0.8)	1318/9970 (13)
Biomarkers of kidney function, median (IQR)							
Serum creatinine, mg/dl ^e	0.77 (0.22)	0.79 (0.22)	0.78 (0.24)	0.75 (0.21)	0.72 (0.21)	0.75 (0.21)	0.82 (0.2)
eGFR, ml/min per 1.73 m ^{2e,f}	92 (21)	100 (23)	95 (25)	102 (17)	104 (13)	100 (16)	100 (20)
Urine albumin, mg/g ^g	4 (10)	1 (6)	3 (8)	4 (8)	2 (3)	1 (2)	2 (6)
Biomarkers of kidney disease^h, <i>N</i> (%)							
Low eGFR ⁱ	114/2197 (5.2)	53/1724 (3.1)	36/1140 (3.2)	55/1768 (3.1)	45/1970 (2.3)	44/1843 (2.4)	347/10,642 (3.3)
Albuminuria ^{g,j}	258/1789 (14)	107/857 (13)	115/1024 (11)	184/1599 (12)	114/1913 (6)	114/1701 (7)	892/8883 (10)
Composite end point ^k	295/1740 (17)	120/856 (14)	142/1021 (14)	222/1596 (14)	144/1902 (8)	151/1700 (9)	1074/8815 (12)
<i>APOL1</i> allele distributions, <i>N</i> (%)							
G0 allele	2999 (67)	2666 (75)	1730 (76)	3108 (88)	3098 (78)	2793 (76)	16,394 (76)
G1 allele	437 (10)	355 (10)	167 (7)	177 (5)	500 (13)	437 (12)	2073 (10)
G2 allele	1070 (24)	529 (15)	389 (17)	257 (7)	368 (9)	458 (12)	3071 (14)
<i>APOL1</i> genotype distributions, <i>N</i> (%)							
0 risk alleles (G0/G0)	997 (44) ^l	1009 (57) ^l	658 (58) ^l	1360 (77) ^l	1210 (61) ^l	1041 (57) ^l	6275 (58) ^l
1 risk allele	1005 (45) ^l	648 (37) ^l	414 (36) ^l	388 (22) ^l	678 (34) ^l	711 (39) ^l	3844 (36) ^l
G0/G1	300 (13)	266 (15)	119 (10)	157 (9)	393 (20)	349 (19)	1584 (15)
G0/G2	705 (31)	382 (22)	295 (26)	231 (13)	285 (14)	362 (20)	2260 (21)
2 risk alleles	251 (11) ^l	118 (7) ^l	71 (6) ^l	23 (1) ^l	95 (5) ^l	92 (5) ^l	650 (6) ^l
G1/G1	14 (0.6)	16 (0.9)	5 (0.4)	4 (0.2)	28 (1.4)	17 (0.9)	84 (0.8)
G1/G2	109 (5)	57 (3)	38 (3)	12 (0.7)	51 (3)	54 (3)	321 (3)
G2/G2	128 (6)	45 (3)	28 (3)	7 (0.4)	16 (0.8)	21 (1)	245 (2)

BMI, body mass index; IQR, interquartile range.

^aAll data are reported as mean (SD) unless stated otherwise. Categories are reported as number (*N*) and percentage; percentages may sum to ± 100 from rounding up.

^bBMI missing data overall $n=456$ (4%): 453 of 456 Agincourt, two of 456 Soweto, and one of 456 Nanoro.

^cDiabetes mellitus missing data overall $n=127$ (1%): 50 of 127 Navrongo, 39 of 127 Agincourt, 16 of 127 Nairobi, ten of 127 Nanoro, seven of 127 Dikgale, and five of 127 Soweto.

^dHIV infection missing data overall $n=799$ (7%): 568 of 799 Soweto, 210 of 799 Nairobi, 19 of 799 Dikgale, and two of 799 Agincourt.

^eSerum creatinine eGFR missing data overall $n=239$ (2%): 91 of 239 Agincourt, 63 of 239 Soweto, 31 of 239 Nanoro, 22 of 239 Navrongo, 21 of 239 Nairobi, and 11 of 239 Dikgale.

^feGFR is calculated using the Chronic Kidney Disease Epidemiology Collaboration (creatinine) equation 2009 without adjusting for the African American coefficient.

^gUrine albumin and albuminuria missing data overall $n=1886$ (18%): 918 of 1886 Soweto, 464 of 1886 Agincourt, 172 of 1886 Nairobi, 143 of 1886 Navrongo, 119 of 1886 Dikgale, and 70 of 1886 Nanoro.

^hFor biomarkers of kidney disease, repeat measures for those with albuminuria or low eGFR were not performed, preventing confirmation of CKD. The category “Biomarkers of kidney disease” was used to define these measures on a single screening.

ⁱLow eGFR is eGFR <60 ml/min per 1.73 m² calculated using the Chronic Kidney Disease Epidemiology Collaboration (creatinine) equation 2009 without adjusting for the African American coefficient.

^jAlbuminuria is random spot urine albumin-creatinine ratio >30 mg/g.

^kThe composite end point is low eGFR and/or albuminuria.

^lCorresponds at main section and in genotype at the total of individual with zero risk allele, one risk allele, and two risk alleles.

site and overall are summarized in Table 1. The mean age of participants was similar across all sites, with some differences in ratios of men to women. Compared with East and West Africa, the three South African sites had the highest prevalence of diabetes mellitus, hypertension, HIV, obesity,

and kidney disease. Overall, *APOL1* allele frequencies were 10% for G1 and 14% for G2. G1 allele frequencies ranged from 5% to 13% across the sites, with the highest frequencies in West Africa. G2 allele frequencies varied widely from 7% to 24%, with the highest frequencies in South Africa. The




Figure 1. | Combined “high-risk” genotype frequencies are highest in West Africa or South Africa and lowest in East Africa, with distinct differences with regard to the G1/G1, G1/G2 and G2/G2 genotypes across regions. *APOL1* genotype (G1/G1, G1/G2, and G2/G2) frequencies (percentages) and combined “high-risk” genotypes (G1/G1, G1/G2, and G2/G2) for each sub-Saharan African region: Nanoro and Navrongo in West Africa; Nairobi in East Africa; and Soweto, Dikgale, and Agincourt study sites in South Africa.

Table 2. Association between *APOL1* and biomarkers of kidney disease by risk allele status (zero, one, or two) and recessive, dominant, and additive models

Biomarkers of Kidney Disease ^a	Low eGFR ^b	Albuminuria ^c	Composite End Point ^d
Frequency of <i>APOL1</i> risk alleles, N (%)			
0	202/6210 (3.3)	507/5259 (10)	627/5225 (12)
1	122/3797 (3.2)	306/3105 (10)	363/3083 (12)
2	23/635 (3.6)	79/519 (15)	84/507 (17)
All	347/10,642 (3.0) ^e	892/8883 (10) ^f	1074/8815 (12)
<i>APOL1</i> models odds ratio (95% confidence interval)			
Sample size, N	9340 ^g	8526 ^g	8486 ^g
Additive	0.78 (0.52 to 1.19)	1.39 (1.09 to 1.76)	1.16 (0.93 to 1.45)
Dominant, 0 versus 1 or 2 risk alleles	0.86 (0.67 to 1.11)	1.12 (0.97 to 1.31)	1.03 (0.90 to 1.19)
Recessive, 0 or 1 versus 2 risk alleles	0.87 (0.52 to 1.47)	1.63 (1.25 to 2.12)	1.37 (1.06 to 1.78)
^a For biomarkers of kidney disease, repeat measures for those with albuminuria or low eGFR were not performed, preventing confirmation of CKD; therefore, we used “biomarkers of kidney disease” to define these measures on a single screening. ^b Low eGFR is eGFR <60 ml/min per 1.73 m ² calculated using the Chronic Kidney Disease Epidemiology Collaboration (creatinine) equation (2009) without adjusting for the African American coefficient. ^c Albuminuria is a random spot urine albumin-creatinine ratio >30 mg/g. ^d The composite end point is low eGFR and/or albuminuria. ^e Missing data n=127 of 10,967 (1%). ^f Missing data n=1886 of 10,967 (17%). ^g Corresponds at main section and in genotype at the total of individual with zero risk allele, one risk allele, and two risk alleles.			

lowest allele frequencies for G1 (5%) and G2 (7%) occurred in East Africa (Table 1). High-risk *APOL1* genotype frequencies were 6% overall, with the lowest in East Africa (1%) and the highest in South Africa (11%). *APOL1* allele and genotype frequencies were highly variable among self-reported ethnic groups (Figure 1, Supplemental Table 3).

***APOL1* Genotypes and Associated Risk for Kidney Disease across All AWI-Gen Studies Sites**

Associations between *APOL1* status (stratified by zero, one, or two risk alleles) and kidney disease risk factors are summarized in Supplemental Table 4. For the study outcomes—low eGFR, albuminuria, and the composite end point—associations with *APOL1* stratified by zero, one, or two risk alleles and using additive, recessive, and dominant models of inheritance are summarized in Table 2. Overall, for low eGFR, there was no association with *APOL1* for any of the models tested. For albuminuria, the association with *APOL1* was significant for the recessive (OR, 1.63; 95% CI, 1.25 to 2.12) and additive models (OR, 1.39; 95% CI, 1.09 to 1.76); for the composite end point, the association was significant for the recessive model (OR, 1.37; 95% CI, 1.06 to 1.78), likely driven by the association with albuminuria. There was no significant heterogeneity ($P=0.78$) when comparing the association with *APOL1* and albuminuria among West, East, and South Africa, and the sample size was likely underpowered to reach statistical significance in East Africa (Supplemental Figure 1, Supplemental Table 5).

The prevalence of obesity was higher in the group with two *APOL1* risk alleles (Supplemental Table 4). As obesity can be causal for albuminuria and we had adjusted our models for BMI, we tested whether the associations with *APOL1* changed without adjusting for BMI and found no difference (Supplemental Table 6). There was no effect of socioeconomic status on the association between *APOL1* and low eGFR, albuminuria, and the composite end point

(Supplemental Table 7). We tested for associations with *APOL1* and low eGFR, albuminuria, and the composite end point stratified by individuals with comorbidity (diabetes mellitus, hypertension, and HIV). *APOL1* was associated with albuminuria in individuals who did not have diabetes mellitus, HT, or HIV, and the interaction term was only significant in the minimally adjusted model (Table 3, Supplemental Tables 9 and 10).

We performed exploratory analyses with different genotype combinations for their association with kidney disease biomarkers. Although there was no association with low eGFR, the association with albuminuria was strongest for G1 homozygotes overall (OR, 3.87; 95% CI, 2.16 to 6.93) and in West Africa (OR, 4.93; 95% CI, 2.15 to 10.26) and South Africa (OR, 3.67; 95% CI, 1.38 to 8.82), and a significant, but smaller effect was seen with G2 homozygotes overall (OR, 1.65; 95% CI, 1.09 to 2.51) and in South Africa (OR, 1.62; 95% CI, 1.01 to 2.51). For the composite end point, only the association with G1 homozygotes remained significant overall (OR, 2.73; 95% CI, 1.51 to 4.96) and in West Africa (OR, 3.45; 95% CI, 1.52 to 7.13). There were no significant associations with the G1/G2 genotype and low eGFR, albuminuria, and the composite end point (Table 4). The differential effect sizes between the two homozygous high-risk genotypes (G1/G1 compared with G2/G2) on the three biomarkers of kidney disease are shown in Supplemental Table 8.

Discussion

We determined population frequencies of *APOL1* alleles and genotypes in multiple ethnic groups from sub-Saharan Africa and investigated their association with biomarkers of kidney disease, leading to several important findings. First, the overall population frequencies of G1 (10%) and G2 (14%) alleles and high-risk genotypes (6%) were lower than reported in African American populations but similar

Table 3. APOL1 associations with albuminuria stratified by diabetes, hypertension, and HIV status

Comorbidity Status	Albuminuria, ^a n/N (%)	Albuminuria ^a Odds Ratio (95% Confidence Interval), Age+Sex-Adjusted Model ^b	Albuminuria ^a Odds Ratio (95% Confidence Interval), Fully Adjusted Model ^c	<i>P</i> _{interaction term} Value, Age+Sex-Adjusted Model ^b
Diabetes				0.31
Absent	745/8259 (9.0)	1.68 (1.28 to 2.20)	1.71 (1.30 to 2.26)	
Present	131/523 (25.0)	1.20 (0.53 to 2.74)	1.06 (0.45 to 2.49)	
Hypertension				0.04
Absent	383/5781 (6.6)	1.94 (1.34 to 2.82)	2.09 (1.43 to 3.05)	
Present	509/3102 (16.4)	1.30 (0.91 to 1.85)	1.33 (0.92 to 1.92)	
HIV				0.40
Negative	655/7508 (8.7)	1.64 (1.22 to 2.19)	1.72 (1.28 to 2.33)	
Positive	202/1129 (17.9)	1.28 (0.74 to 2.23)	1.36 (0.78 to 2.40)	

^aAlbuminuria is random spot urine albumin-creatinine ratio >30 mg/g.
^bLogistic mixed models were adjusted for age and sex as fixed variables.
^cLogistic mixed models were adjusted for site as a random variable and age, sex, body mass index, diabetes mellitus status, hypertension status, and HIV status as fixed variables. Some participants were excluded from the fully adjusted model due to missing data.

to a recent study from Nigeria, West Africa (high-risk genotype: 6%) (17). We observed distinct differences in population frequencies of *APOL1* alleles and high-risk genotypes when comparing West, East, and South African regions and among ethnic groups within each region, with the lowest frequencies among East African ethnic groups.

Second, *APOL1* high-risk genotypes were strongly associated with albuminuria but not with low eGFR. The association with albuminuria is consistent with continental African

(29) and African American studies (7,40,41). The absence of association with low eGFR might be explained by the use of GFR estimating equations that have limited validation in sub-Saharan Africa, including little difference between the CKD-EPI (creatinine) 2009 and 2021 equations (Supplemental Appendix 1), and laboratory use of the Jaffe rather than the enzymatic assay for creatinine—both of which tend toward overestimating eGFR. It is also plausible that a survival or frailty bias resulting from the near absence of

Table 4. Association between APOL1 genotype and biomarkers of kidney disease stratified by region

Genotype ^a	All	East Africa	South Africa	West Africa
Low eGFR^b				
G0/G1	1.03 (0.73 to 1.47)	1.00 (0.29 to 2.63)	1.06 (0.61 to 1.76)	1.01 (0.58 to 1.69)
G0/G2	0.76 (0.55 to 1.05)	0.77 (0.26 to 1.88)	0.82 (0.53 to 1.25)	0.74 (0.37 to 1.34)
G1/G1	0.52 (0.07 to 3.83)	NA ^c	0 (0 to 35,130.28)	0.93 (0.05 to 4.47)
G1/G2	0.81 (0.39 to 1.69)	4.0 (0.2 to 26.4)	1.13 (0.46 to 2.38)	NA ^c
G2/G2	0.89 (0.41 to 1.94)	15.09 (0.69 to 125.23)	0.99 (0.38 to 2.18)	NA ^c
Albuminuria^d				
G0/G1	0.97 (0.77 to 1.22)	0.79 (0.38 to 1.48)	0.82 (0.58 to 1.14)	1.3 (0.9 to 1.8)
G0/G2	1.09 (0.9 to 1.31)	1.21 (0.73 to 1.95)	1.06 (0.84 to 1.35)	1.1 (0.7 to 1.6)
G1/G1	3.87 (2.16 to 6.93) ^e	NA ^c	3.67 (1.38 to 8.82) ^e	4.93 (2.15 to 10.26) ^e
G1/G2	1.24 (0.83 to 1.87)	3.24 (0.16 to 23.79)	1.20 (0.72 to 1.92)	1.28 (0.52 to 2.66)
G2/G2	1.65 (1.09 to 2.51) ^e	4.63 (0.62 to 23.40)	1.62 (1.01 to 2.51) ^e	1.09 (0.17 to 3.71)
Composite end point^f				
G0/G1	0.97 (0.79 to 1.19)	0.84 (0.44 to 1.50)	0.83 (0.61 to 1.13)	1.22 (0.89 to 1.65)
G0/G2	0.99 (0.83 to 1.17)	1.12 (0.69 to 1.77)	0.98 (0.78 to 1.22)	0.95 (0.67 to 1.34)
G1/G1	2.73 (1.51 to 4.96) ^e	NA ^c	2.47 (0.87 to 6.18)	3.45 (1.52 to 7.13) ^e
G1/G2	1.06 (0.72 to 1.57)	2.56 (0.12 to 19.26)	1.11 (0.69 to 1.73)	0.89 (0.37 to 1.83)
G2/G2	1.40 (0.93 to 2.11)	4.32 (0.59 to 21.73)	1.41 (0.89 to 2.17)	0.74 (0.12 to 2.52)

The associations between *APOL1* genotype and biomarkers of kidney disease are presented as odds ratios and 95% confidence intervals. For biomarkers of kidney disease, repeat measures for those with low eGFR or albuminuria were not performed, thus preventing confirmation of CKD; therefore, we used “biomarkers of kidney disease” to define these measures on a single screening. Linear and logistic mixed models were adjusted for site as a random variable and age, sex, body mass index, diabetes mellitus status, hypertension status, and HIV status as fixed variables. NA, not applicable.

^a*APOL1* genotypes G0/G1, G0/G2, G1/G1, G1/G2, and G2/G2 were compared with G0/G0 as the reference.

^bLow eGFR is eGFR <60 ml/min per 1.73 m² calculated using the Chronic Kidney Disease Epidemiology Collaboration (creatinine) equation 2009 without adjusting for the African American coefficient.

^cSample size was too small to perform the statistical test.

^dAlbuminuria is random spot urine albumin-creatinine ratio >3.0 mg/g.

^eCorresponds at main section and in genotype at the total of individual with zero risk allele, one risk allele, and two risk alleles.

^fThe composite end point is low eGFR and/or albuminuria.

access to KRT for those with kidney failure accounts for the lack of association with low eGFR because those with progressive kidney disease due to *APOL1* might be under-represented in our sample. Finally, the population-based study design did not select for groups with severe or progressive kidney disease.

Third, regarding kidney disease risk, we showed an association with obesity and two *APOL1* risk alleles but no demonstrable effect of obesity on the association of *APOL1* with albuminuria. Our findings corroborate those of a similar-sized study of 11,930 African American participants, where each risk allele was associated with greater obesity odds (1.13-fold) (42), but this association has not been demonstrated in other large cohorts, such as the Coronary Artery Risk Development in Young Adults and the Atherosclerosis Risk in Communities studies (19,43). There was no effect of socioeconomic status on the association between *APOL1* and albuminuria. Although this finding is similar to those observed in African American participants (19,43), the assets-based score differs from scoring systems used in high-income settings, and quintile differences may be difficult to detect when the overall status of African participants is one of widespread poverty. There was a significant interaction when testing the *APOL1* association with albuminuria for the minimally adjusted model between participants with and without hypertension, but no significant interactions between established risk factors (diabetes, hypertension, and HIV) and *APOL1* risk alleles in the fully adjusted model. This lack of interaction could be due to the limited sample size of subgroups of individuals with HIV and diabetes.

Lastly, differential effects of G1 and G2 homozygous genotypes on albuminuria are similar to those reported in the Dallas and Jackson Heart studies, but contradict other studies in which high-risk genotypes have near-equivalent genetic associations in the homozygous or compound heterozygous states (7,24). The absence of association with *APOL1* G1/G2 and albuminuria has not been previously described and is difficult to explain, especially because G1/G2 represents the largest group of high-risk genotypes. Studies have shown functional differences between G1 and G2, with their proteins showing differential binding affinities to the trypanosome-encoded serum resistance-associated protein and associations with susceptibility to *T.b. rhodiense* and *T.b. gambiense* infection and disease (27). Proposed mechanisms of *APOL1*-induced cytotoxicity in kidney tissue using *in vitro* and mouse models include enhanced local *APOL1* expression and reduced degradation, increased cation transport, downstream intracellular activation of the inflammasome, defects in autophagy, and mitochondrial dysfunction. However, none of these demonstrate differential effects among high-risk genotypes (16).

Our study has important strengths and limitations. Strengths lie in the well-characterized cohort; the large population-based sampling frame; inclusion of multiple ethnic groups from countries at different stages of the health and epidemiologic transition in West, East, and South Africa; and the availability of GFR estimates and albuminuria as biomarkers of kidney disease. The cross-sectional study design is a limitation as single measurements of eGFR and albuminuria preclude CKD diagnosis. Although albuminuria is a marker for CKD progression, it can also be

reversible or regress with time, which is particularly relevant in sub-Saharan Africa where acute, chronic, and endemic infectious disease burdens are high, as are heavy metal and toxin exposures. Without longitudinal follow-up, the effect of high-risk *APOL1* genotypes on incident albuminuria, incident CKD, and CKD progression cannot be evaluated. As our analyses were exploratory and sample sizes were relatively small for some subgroup comparisons, our findings should be considered hypothesis generating and will hopefully inform future work for validation.

In conclusion, population frequencies of *APOL1* G1 and G2 alleles and high-risk genotypes differed between and within West and South Africa and were almost absent from East Africa. *APOL1* kidney risk variants were associated with albuminuria but not eGFR. We noted differential effects of homozygous G1 and G2 genotypes on albuminuria, which might result from genetic or environmental modifiers. Compared with findings in African American populations, the relatively low frequencies and variable penetrance of *APOL1* high-risk genotypes, regional differences in the association with albuminuria, and absence of association with low eGFR may reflect the vast genetic diversity of African populations, differences in gene-environment interactions or genetic modifiers in resident African populations, and the absence of European admixture. Further studies in sub-Saharan Africa with data from different regions and ethnic groups are required to confirm and refine the effect of *APOL1* risk variants on kidney phenotypes, and longitudinal studies are needed to assess the role of *APOL1* and CKD incidence and progression.

Disclosures

M. Ramsay reports consultancy agreements with Roche/Genentech (The Health Equity and Population Science). M. Ramsay is a South African Research Chair in Genomics and Bioinformatics of African Populations hosted by the University of the Witwatersrand, funded by the Department of Science and Technology, and administered by the National Research Foundation. All remaining authors have nothing to disclose.

Funding

The AWI-Gen Collaborative Centre is funded by National Human Genome Research Institute award U54HG006938 and its supplements as part of the H3Africa Consortium. Additional funding came from Department of Science and Technology, South Africa award DST/CON 0056/2014. The project has been supported in part by NIH Clinical Center contract 75N91019D00024.

Acknowledgments

We thank all AWI-Gen participants and acknowledge our field workers, laboratory scientists, administrators, data manager, and other staff who contributed to data collection at each study site (Supplemental Appendix 2). In particular, we acknowledge Cassandra Soo, Freedom Mukomana, and Scott Hazelhurst from the Sydney Brenner Institute for Molecular Bioscience; Stephen Tollman; Kathleen Kahn (Agincourt Principal Investigator [PI]); Marianne Alberts (Dikgale PI, deceased in 2020); Catherine Kyobutungi (Nairobi PI); Halidou Tinto (Nanoro PI); Abraham R. Oduro (Navrongo PI); and Shane Norris (Soweto PI).

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

The content of this publication does not necessarily reflect the view or policy of the funders, nor does mention of trade names, commercial products, or organizations imply endorsement by the US or South African governments.

Author Contributions

J.-T. Brandenburg, J. Fabian, and M. Ramsay conceptualized the study; J.-T. Brandenburg, J. Fabian, M.A. Govender, M. Ramsay, and C.A. Winkler were responsible for formal analysis; J.-T. Brandenburg and M.A. Govender wrote the original draft; and G. Agongo, P.R. Boua, J.-T. Brandenburg, J. Fabian, M.A. Govender, M. Ramsay, and C.A. Winkler reviewed and edited the manuscript.

Data Sharing Statement

The phenotype and genotype data are available on request to the H3Africa Data and Biospecimen Access Committee (European Genome-Phenome Archive accession nos. EGAD00001006425 and EGAD00001001996, respectively).

Supplemental Material

This article contains the following supplemental material online at <http://cjasn.asnjournals.org/lookup/suppl/doi:10.2215/CJN.14321121/-/DCSupplemental>.

Supplemental Appendix 1. Comparison of models between eGFR using CKD-EPI 2009 and CKD-EPI 2021.

Supplemental Appendix 2. The H3Africa Consortium AWI-Gen Study.

Supplemental Table 1. *APOL1* haplotypes (referred to as alleles G0, G1, and G2 for ease of comparison with other literature) on the basis of three polymorphic loci on chromosome 22.

Supplemental Table 2. Quality of imputation for the three variant positions and *APOL1* G0, G1, and G2 risk alleles on chromosome 22.

Supplemental Table 3. *APOL1* alleles and high-risk genotypes using recessive models by ethnicity (data used for Figure 1).

Supplemental Table 4. *APOL1* risk alleles and associated risk factors for kidney disease across the combined dataset of the Africa Wits–International Network for the Demographic Evaluation of Populations and Their Health Partnership for Genomic Studies.

Supplemental Table 5. *APOL1* associations with albuminuria across the Africa Wits–International Network for the Demographic Evaluation of Populations and Their Health Partnership for Genomic Studies study sites.

Supplemental Table 6. Association (OR; 95% CI) between *APOL1* using additive, dominant, and recessive models and albuminuria alone or as a composite end point.

Supplemental Table 7. Association (OR; 95% CI) between *APOL1* using additive, dominant, and recessive models and albuminuria, low eGFR, and the composite end point.

Supplemental Table 8. Effect of genotypes (G1/G1 compared with G2/G2) on biomarkers of kidney disease.

Supplemental Table 9. *APOL1* associations with low eGFR stratified by diabetes, hypertension, and HIV status.

Supplemental Table 10. *APOL1* associations with the composite end point stratified by diabetes, hypertension, and HIV status.

Supplemental Figure 1. Forest plot: association between high-risk *APOL1* genotypes (OR; 95% CI) and albuminuria by region.

References

- Ashuntantang G, Osafo C, Olowu WA, Arogundade F, Niang A, Porter J, Naicker S, Luyckx VA: Outcomes in adults and

- children with end-stage kidney disease requiring dialysis in sub-Saharan Africa: A systematic review. *Lancet Glob Health* 5: e408–e417, 2017
- Kaze AD, Ilori T, Jaar BG, Echouffo-Tcheugui JB: Burden of chronic kidney disease on the African continent: A systematic review and meta-analysis. *BMC Nephrol* 19: 125, 2018
- Kopp JB, Smith MW, Nelson GW, Johnson RC, Freedman BI, Bowden DW, Oleksyk T, McKenzie LM, Kajiyama H, Ahuja TS, Berns JS, Briggs W, Cho ME, Dart RA, Kimmel PL, Korbet SM, Michel DM, Mokrzycki MH, Schelling JR, Simon E, Trachtman H, Vlahov D, Winkler CA: MYH9 is a major-effect risk gene for focal segmental glomerulosclerosis. *Nat Genet* 40: 1175–1184, 2008
- Kao WH, Klag MJ, Meoni LA, Reich D, Berthier-Schaad Y, Li M, Coresh J, Patterson N, Tandon A, Powe NR, Fink NE, Sadler JH, Weir MR, Abboud HE, Adler SG, Divers J, Iyengar SK, Freedman BI, Kimmel PL, Knowler WC, Kohn OF, Kramp K, Leehey DJ, Nicholas SB, Pahl MV, Schelling JR, Sedor JR, Thornley-Brown D, Winkler CA, Smith MW, Parekh RS; Family Investigation of Nephropathy and Diabetes Research Group: MYH9 is associated with nondiabetic end-stage renal disease in African Americans. *Nat Genet* 40: 1185–1192, 2008
- Genovese G, Friedman DJ, Ross MD, Lecordier L, Uzureau P, Freedman BI, Bowden DW, Langefeld CD, Oleksyk TK, Uscinski Knob AL, Bernhardt AJ, Hicks PJ, Nelson GW, Vanhollebeke B, Winkler CA, Kopp JB, Pays E, Pollak MR: Association of trypanolytic *ApoL1* variants with kidney disease in African Americans. *Science* 329: 841–845, 2010
- Kopp JB, Nelson GW, Sampath K, Johnson RC, Genovese G, An P, Friedman D, Briggs W, Dart R, Korbet S, Mokrzycki MH, Kimmel PL, Limou S, Ahuja TS, Berns JS, Fryc J, Simon EE, Smith MC, Trachtman H, Michel DM, Schelling JR, Vlahov D, Pollak M, Winkler CA: *APOL1* genetic variants in focal segmental glomerulosclerosis and HIV-associated nephropathy. *J Am Soc Nephrol* 22: 2129–2137, 2011
- Friedman DJ, Kozlitina J, Genovese G, Jog P, Pollak MR: Population-based risk assessment of *APOL1* on renal disease. *J Am Soc Nephrol* 22: 2098–2105, 2011
- Madhavan SM, O'Toole JF, Konieczkowski M, Ganesan S, Bruggeman LA, Sedor JR: *APOL1* localization in normal kidney and nondiabetic kidney disease. *J Am Soc Nephrol* 22: 2119–2128, 2011
- Tzur S, Rosset S, Shemer R, Yudkovsky G, Selig S, Tarekegn A, Bekele E, Bradman N, Wasser WG, Behar DM, Skorecki K: Missense mutations in the *APOL1* gene are highly associated with end stage kidney disease risk previously attributed to the MYH9 gene. *Hum Genet* 128: 345–350, 2010
- Pays E, Vanhollebeke B, Uzureau P, Lecordier L, Pérez-Morga D: The molecular arms race between African trypanosomes and humans. *Nat Rev Microbiol* 12: 575–584, 2014
- Papeta N, Kiryluk K, Patel A, Sterken R, Kacak N, Snyder HJ, Imus PH, Mhatre AN, Lawani AK, Julian BA, Wyatt RJ, Novak J, Wyatt CM, Ross MJ, Winston JA, Klotman ME, Cohen DJ, Appel GB, D'Agati VD, Klotman PE, Gharavi AG: *APOL1* variants increase risk for FSGS and HIVAN but not IgA nephropathy. *J Am Soc Nephrol* 22: 1991–1996, 2011
- Larsen CP, Beggs ML, Saeed M, Walker PD: Apolipoprotein L1 risk variants associate with systemic lupus erythematosus-associated collapsing glomerulopathy. *J Am Soc Nephrol* 24: 722–725, 2013
- Nichols B, Jog P, Lee JH, Blackler D, Wilmot M, D'Agati V, Markowitz G, Kopp JB, Alper SL, Pollak MR, Friedman DJ: Innate immunity pathways regulate the nephropathy gene Apolipoprotein L1. *Kidney Int* 87: 332–342, 2015
- Velez JCQ, Caza T, Larsen CP: COVAN is the new HIVAN: The re-emergence of collapsing glomerulopathy with COVID-19. *Nat Rev Nephrol* 16: 565–567, 2020
- Kasembeli AN, Duarte R, Ramsay M, Mosiane P, Dickens C, Dix-Peek T, Limou S, Sezgin E, Nelson GW, Fogo AB, Goetsch S, Kopp JB, Winkler CA, Naicker S: *APOL1* risk variants are strongly associated with HIV-associated nephropathy in black South Africans. *J Am Soc Nephrol* 26: 2882–2890, 2015
- Freedman BI, Kopp JB, Sampson MG, Susztak K: *APOL1* at 10 years: Progress and next steps. *Kidney Int* 99: 1296–1302, 2021

17. Limou S, Nelson GW, Kopp JB, Winkler CA: APOL1 kidney risk alleles: Population genetics and disease associations. *Adv Chronic Kidney Dis* 21: 426–433, 2014 10.1053/j.ackd.2014.06.005
18. Kruzell-Davila E, Wasser WG, Aviram S, Skorecki K: APOL1 nephropathy: From gene to mechanisms of kidney injury. *Nephrol Dial Transplant* 31: 349–358, 2016
19. Grams ME, Rebholz CM, Chen Y, Rawlings AM, Estrella MM, Selvin E, Appel LJ, Tin A, Coresh J: Race, APOL1 risk, and eGFR decline in the general population. *J Am Soc Nephrol* 27: 2842–2850, 2016
20. Palmer ND, Freedman BI: APOL1 and progression of nondiabetic nephropathy. *J Am Soc Nephrol* 24: 1344–1346, 2013
21. Foster MC, Coresh J, Fornage M, Astor BC, Grams M, Franceschini N, Boerwinkle E, Parekh RS, Kao WH: APOL1 variants associate with increased risk of CKD among African Americans. *J Am Soc Nephrol* 24: 1484–1491, 2013
22. Lipkowitz MS, Freedman BI, Langefeld CD, Comeau ME, Bowden DW, Kao WH, Astor BC, Bottinger EP, Iyengar SK, Klotman PE, Freedman RG, Zhang W, Parekh RS, Choi MJ, Nelson GW, Winkler CA, Kopp JB; SK Investigators: Apolipoprotein L1 gene variants associate with hypertension-attributed nephropathy and the rate of kidney function decline in African Americans. *Kidney Int* 83: 114–120, 2013
23. Parsa A, Kao WH, Xie D, Astor BC, Li M, Hsu CY, Feldman HI, Parekh RS, Kusek JW, Greene TH, Fink JC, Anderson AH, Choi MJ, Wright Jr. JT, Lash JP, Freedman BI, Ojo A, Winkler CA, Raj DS, Kopp JB, He J, Jensvold NG, Tao K, Lipkowitz MS, Appel LJ; AASK Study Investigators; CRIC Study Investigators: APOL1 risk variants, race, and progression of chronic kidney disease. *N Engl J Med* 369: 2183–2196, 2013
24. Ito K, Bick AG, Flannick J, Friedman DJ, Genovese G, Parfenov MG, Depalma SR, Gupta N, Gabriel SB, Taylor Jr. HA, Fox ER, Newton-Cheh C, Kathiresan S, Hirschhorn JN, Altshuler DM, Pollak MR, Wilson JG, Seidman JG, Seidman C: Increased burden of cardiovascular disease in carriers of APOL1 genetic variants. *Circ Res* 114: 845–850, 2014
25. Kanji Z, Powe CE, Wenger JB, Huang C, Ankers E, Sullivan DA, Collerone G, Powe NR, Tonelli M, Bhan I, Bernhardt AJ, Dibartolo S, Friedman D, Genovese G, Pollak MR, Thadhani R: Genetic variation in APOL1 associates with younger age at hemodialysis initiation. *J Am Soc Nephrol* 22: 2091–2097, 2011
26. Tzur S, Rosset S, Skorecki K, Wasser WG: APOL1 allelic variants are associated with lower age of dialysis initiation and thereby increased dialysis vintage in African and Hispanic Americans with non-diabetic end-stage kidney disease. *Nephrol Dial Transplant* 27: 1498–1505, 2012
27. Friedman DJ, Pollak MR: APOL1 and kidney disease: From genetics to biology. *Annu Rev Physiol* 82: 323–342, 2020
28. Behar DM, Kedem E, Rosset S, Haileselassie Y, Tzur S, Kra-Oz Z, Wasser WG, Shenhar Y, Shahar E, Hassoun G, Maor C, Wolday D, Pollack S, Skorecki K: Absence of APOL1 risk variants protects against HIV-associated nephropathy in the Ethiopian population. *Am J Nephrol* 34: 452–459, 2011
29. Sumaili EK, Shemer R, Kruzell-Davila E, Cohen EP, Mutantun PN, Bukabau JB, Makulo JRR, Mokoli VM, Luse JL, Pakasa NM, Cavalier E, Wumba RD, Reiner-Benaïm A, Boner G, Lifschitz M, Nseka NM, Skorecki K, Wasser WG: G1 is the major APOL1 risk allele for hypertension-attributed nephropathy in Central Africa. *Clin Kidney J* 12: 188–195, 2018
30. Ulasi II, Tzur S, Wasser WG, Shemer R, Kruzell E, Feigin E, Ijoma CK, Onodugo OD, Okoye JU, Arodiwe EB, Ifeunandu NA, Chukwuka CJ, Onyedum CC, Ijoma UN, Nna E, Onuigbo M, Rosset S, Skorecki K: High population frequencies of APOL1 risk variants are associated with increased prevalence of non-diabetic chronic kidney disease in the Igbo people from south-eastern Nigeria. *Nephron Clin Pract* 123: 123–128, 2013
31. Wudil UJ, Aliyu MH, Prigmore HL, Ingles DJ, Ahonkhai AA, Musa BM, Muhammad H, Sani MU, Nalado AM, Abdu A, Abdussalam K, Shepherd BE, Dankishiya FS, Burgner AM, Iki-zler TA, Wyatt CM, Kopp JB, Kimmel PL, Winkler CA, Wester CW: Apolipoprotein-1 risk variants and associated kidney phenotypes in an adult HIV cohort in Nigeria. *Kidney Int* 100: 146–154, 2021
32. Ramsay M, Crowther N, Tambo E, Agongo G, Baloyi V, Dikotopé S, Gómez-Olivé X, Jaff N, Sorgho H, Wagner R, Khayeka-Wandabwa C, Choudhury A, Hazelhurst S, Kahn K, Lombard Z, Mukomana F, Soo C, Soodyall H, Wade A, Afolabi S, Agorinya I, Amenga-Etego L, Ali SA, Bognini JD, Boua RP, Debpuur C, Diallo S, Fato E, Kazienga A, Konkobo SZ, Kouraogo PM, Mashinya F, Micklesfield L, Nakanabo-Diallo S, Njamwea B, Nonterah E, Ouedraogo S, Pillay V, Somande AM, Tindana P, Twine R, Alberts M, Kyobutungi C, Norris SA, Oduro AR, Tinto H, Tollman S, Sankoh O: H3Africa AWI-Gen Collaborative Centre: A resource to study the interplay between genomic and environmental risk factors for cardiometabolic diseases in four sub-Saharan African countries. *Glob Health Epidemiol Genom* 1: e20, 2016
33. Ali SA, Soo C, Agongo G, Alberts M, Amenga-Etego L, Boua RP, Choudhury A, Crowther NJ, Depuur C, Gómez-Olivé FX, Guiraud I, Haregu TN, Hazelhurst S, Kahn K, Khayeka-Wandabwa C, Kyobutungi C, Lombard Z, Mashinya F, Micklesfield L, Mohamed SF, Mukomana F, Nakanabo-Diallo S, Natama HM, Ngomi N, Nonterah EA, Norris SA, Oduro AR, Somé AM, Sorgho H, Tindana P, Tinto H, Tollman S, Twine R, Wade A, Sankoh O, Ramsay M: Genomic and environmental risk factors for cardiometabolic diseases in Africa: Methods used for phase 1 of the AWI-Gen population cross-sectional study. *Glob Health Action* 11[Suppl2]: 1507133, 2018
34. Levey AS, Stevens LA, Schmid CH, Zhang YL, Castro 3rd AF, Feldman HI, Kusek JW, Eggers P, Van Lente F, Greene T, Coresh J; CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration): A new equation to estimate glomerular filtration rate. *Ann Intern Med* 150: 604–612, 2009
35. McCarthy S, Das S, Kretschmar W, Delaneau O, Wood AR, Teumer A, Kang HM, Fuchsberger C, Danecek P, Sharp K, Luo Y, Sidore C, Kwong A, Timpson N, Koskinen S, Vrieze S, Scott LJ, Zhang H, Mahajan A, Veldink J, Peters U, Pato C, van Duijn CM, Gillies CE, Gandin I, Mezzavilla M, Gilly A, Cocca M, Traglia M, Angius A, Barrett JC, Boomsma D, Branham K, Breen G, Brummett CM, Busonero F, Campbell H, Chan A, Chen S, Chew E, Collins FS, Corbin LJ, Smith GD, Dedoussis G, Dorr M, Farmaki AE, Ferrucci L, Forer L, Fraser RM, Gabriel S, Levy S, Groop L, Harrison T, Hattersley A, Holmen OL, Hveem K, Kretzler M, Lee JC, McGue M, Meitinger T, Melzer D, Min JL, Mohlke KL, Vincent JB, Nauck M, Nickerson D, Palotie A, Pato M, Pirastu N, McInnis M, Richards JB, Sala C, Salomaa V, Schlessinger D, Schoenherr S, Slagboom PE, Small K, Spector T, Stambolian D, Tuke M, Tuomilehto J, Van den Berg LH, Van Rheenen W, Volker U, Wijmenga C, Toniolo D, Zeggiani E, Gasparini P, Sampson MG, Wilson JF, Frayling T, de Bakker PI, Swertz MA, McCarroll S, Kooperberg C, Dekker A, Altshuler D, Willer C, Iacono W, Ripatti S, Soranzo N, Walter K, Swaroop A, Cucca F, Anderson CA, Myers RM, Boehnke M, McCarthy MI, Durbin R; Haplotype Reference Consortium: A reference panel of 64,976 haplotypes for genotype imputation. *Nat Genet* 48: 1279–1283, 2016
36. Choudhury A, Aron S, Botigué LR, Sengupta D, Botha G, Bensellak T, Wells G, Kumuthini J, Shriner D, Fakim YJ, Ghoorah AW, Dareng E, Odia T, Falola O, Adebisi Y, Hazelhurst S, Mazandu G, Nyangiri OA, Mbiyavanga M, Benkahla A, Kassim SK, Mulder N, Adebamowo SN, Chimusa ER, Muzny D, Metcalf G, Gibbs RA, Rotimi C, Ramsay M, Adeyemo AA, Lom-bard Z, Hanchard NA; TrypanoGEN Research Group; H3Africa Consortium: High-depth African genomes inform human migration and health. *Nature* 586: 741–748, 2020
37. Auton A, Brooks LD, Durbin RM, Garrison EP, Kang HM, Korbel JO, Marchini JL, McCarthy S, McVean GA, Abecasis GR; 1000 Genomes Project Consortium: A global reference for human genetic variation. *Nature* 526: 68–74, 2015
38. Gurdasani D, Carstensen T, Tekola-Ayele F, Pagani L, Tachmazidou I, Hatzikotoulas K, Karthikeyan S, Iles L, Pollard MO, Choudhury A, Ritchie GR, Xue Y, Asimit J, Nsubuga RN, Young EH, Pomilla C, Kivinen K, Rockett K, Kamali A, Doumaty AP, Asiki G, Seeley J, Sisay-Joof F, Jallow M, Tollman S, Mekonnen E, Ekong R, Oljira T, Bradman N, Bojang K, Ramsay M, Adeyemo A, Bekele E, Motala A, Norris SA, Pirie F, Kaleebu P, Kwiatkowski D, Tyler-Smith C, Rotimi C, Zeggini E, Sandhu MS: The African Genome Variation Project shapes medical genetics in Africa. *Nature* 517: 327–332, 2015

39. Kuznetsova A, Brockhoff PB, Christensen RH: lmerTest package: Tests in linear mixed effects models. *J Stat Softw* 82: 1–26, 2017
40. Peralta CA, Bibbins-Domingo K, Vittinghoff E, Lin F, Fornage M, Kopp JB, Winkler CA: APOL1 genotype and race differences in incident albuminuria and renal function decline. *J Am Soc Nephrol* 27: 887–893, 2016
41. Chen TK, Tin A, Peralta CA, Appel LJ, Choi MJ, Lipkowitz MS, Winkler CA, Estrella MM: APOL1 risk variants, incident proteinuria, and subsequent eGFR decline in blacks with hypertension-attributed CKD. *Clin J Am Soc Nephrol* 12: 1771–1777, 2017
42. Nadkarni GN, Fei K, Galarneau G, Gao Y, Wilson JG, Cooper R, Madden EB, Denny JC, Richardson LD, Pollak M, Loos RJF, Horowitz CR: APOL1 renal risk variants are associated with obesity and body composition in African ancestry adults: An observational genotype-phenotype association study. *Medicine (Baltimore)* 100: e27785, 2021
43. Chen TK, Estrella MM, Vittinghoff E, Lin F, Gutierrez OM, Kramer H, Lewis CE, Kopp JB, Allen NB, Winkler CA, Bibbins-Domingo KB, Peralta CA: APOL1 genetic variants are not associated with longitudinal blood pressure in young black adults. *Kidney Int* 92: 964–971, 2017

Received: November 2, 2021 **Accepted:** March 30, 2022

J.F. and M.R. contributed equally to this work.

Published online ahead of print. Publication date available at www.cjasn.org.

AFFILIATIONS

¹Sydney Brenner Institute for Molecular Bioscience, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

²Division of Human Genetics, National Health Laboratory Service and School of Pathology, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

³Molecular Genetic Epidemiology Section, Basic Research Laboratory, Frederick National Laboratory for Cancer Research, Frederick, Maryland

⁴Clinical Research Unit of Nanoro, Institut de Recherche en Sciences de la Santé–DRCO, Nanoro, Burkina Faso

⁵Navrongo Health Research Centre, Ghana Health Service, Navrongo, Ghana

⁶Department of Biochemistry and Forensic Sciences, School of Chemical and Biochemical Sciences, C.K. Tedam University of Technology and Applied Sciences, Navrongo, Ghana

⁷Wits Donald Gordon Medical Centre, School of Clinical Medicine, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa

⁸Medical Research Council/Wits University Rural Public Health and Health Transitions Research Unit (Agincourt), School of Public Health, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa